

### SOP 3.8.1 DNA Extraction from Cells in Urine

**SOP Number:** 3.8.1  
**Version Number** 1.0

	Name	Title	Date
Author			
Authoriser			

Effective Date	
Version Number	

#### Purpose

This SOP describes the procedure for DNA extraction from cells in urine.

#### Responsibility

It is the responsibility of the research laboratory personnel carrying out this procedure to ensure that all steps are completed both competently and safely.

#### Equipment/reagent requirements

- DNA extraction buffer
- Proteinase K
- Protein precipitation solution
- Isopropanol
- Ethanol
- DNA hydration buffer
- An incubator/water-bath capable of 30°C
- A refrigerated centrifuge capable of 12,000g
- Vortex mixer
- -80°C Freezer
- A spectrophotometer capable of reading absorbance at 260 and 280nm / Nanodrop

#### Procedure

1. Resuspend the cell pellet from step 4 in SOP 3.8 in 180µL DNA extraction buffer (0.1M EDTA, 0.2M NaCl, 0.05M Tris-HCl, pH 8.0, 0.5% SDS, 50µg/mL DNase free RNase).
2. Add proteinase K to a final concentration of 100µg/mL and gently swirl the tube to mix the components. Incubate the tube at 37°C for at least 3 hours, preferably overnight, with gentle agitation. This can be achieved using a shaking water bath or by occasional swirling by hand. The solution should be reasonably clear and viscous at the end of the incubation.

3. Add 100 $\mu$ L of ammonium acetate protein precipitation solution and vortex vigorously for 20 sec at high speed.
4. Centrifuge at 12,000g for 5 min. The precipitated proteins should form a tight, dark brown pellet. If the protein pellet is not tight, incubate on ice for 5 min and repeat the centrifugation step.
5. Dispense 300 $\mu$ L isopropanol into a clean 1.5mL eppendorf tube and add the supernatant from the previous step. It is important to ensure that the protein pellet is not dislodged during pouring.
6. Mix by inverting gently 50 times until the DNA is visible as threads or a clump. Centrifuge at 12,000g for 3 min. Discard supernatant carefully. Drain the tube on a clean piece of absorbent paper, taking care that the pellet remains in the tube.
7. Wash the DNA, by adding 300 $\mu$ L of ethanol (70% v/v) and vortex at medium speed for 5 sec to dislodge the pellet from the bottom of the tube. Centrifuge at 12,000g for 3 min and drain to remove ethanol, ensure that the DNA pellet remains in the tube.
8. Add 50 $\mu$ L of DNA hydration buffer (TE buffer) and vortex for 5 sec at medium speed to mix. Incubate at 65°C for 1 hour to dissolve DNA. Incubate at RT overnight with gentle shaking. Ensure tube cap is tightly closed to avoid leakage. Centrifuge samples briefly.
9. The absorbance of the DNA at 260nm and 280nm should be measured using quartz cuvetts or the nanodroplet method to assess purity. A 260/280 ratio between 1.8 and 2.0 is desirable. A 260/280 ratio greater than 2.0 may indicate solvent contamination and a ratio less than 1.8 may indicate protein contamination. DNA concentration can be measured using absorbance at 260nm with an  $A_{260}$  of 1.0 in a 1-cm light path being equivalent to a DNA concentration of 50 $\mu$ g/mL / nanodrop. The DNA sample is aliquoted into cooled cryostorage tubes and stored at -80°C.
10. An aliquot of the DNA from a representative sample from each batch may be analyzed by electrophoresis through a 0.3% agarose gel. The prepared DNA is normally at least 100kbp and preferably exceeds 200kbp.

**Note:** There are a number of commercially available kits based on the salting out methods from different manufacturers. When using these methods follow the manufacturer's instructions as outlined in the information for use/ package insert included with the kit. The method used for DNA extraction should be recorded in the study specific documentation or data management system.

**Change History**

<b>SOP Number</b>	<b>Effective Date</b>	<b>Significant Change</b>	<b>Previous SOP No.</b>